
Content

Editorial Page

Functional Evaluation for Planning Exercise Programs in Patients with Chronic Diseases ii

Original Articles

The Effect of Paretic Side Quadriceps Resistive Exercise on Weight-Bearing Asymmetry in Standing Position and Lean Muscle Mass of Subacute Stroke Patients 77

Yenny SEI, Arisanti F and Prabowo T

Effects of Home-based Exercise Program Using Thai-style Braided Rubber Rope on Blood Pressure, Muscle Strength and Quality of Life in Patients on Continuous Ambulatory Peritoneal Dialysis 81

Aramrussameekul W and Changsirikunchai S

Prevalence and Potential Associated Factors of Poststroke Depression 85

Dajpratham P, Pukrittayakamee P, Atsariyasing W, Wannarit K, Boonhong J, Pongpirul K, Suesuwan O and Chatdharmmaluck S

Prevalence and Risk Factors of Serious Arrhythmia during 6-minute Walk Test in Phase II Cardiac Rehabilitation 90

Kwanchuay P and Chitkresorn P

Relationship Between 6-minute Walk Distance and Risk of Hospitalization from Exacerbation of COPD in One Year at Trang Hospital 94

Poosiri S

A Trigger Point Injection with 1% versus 2% Lidocaine for Treatment of Myofascial Pain Syndrome at Neck and Upper Back: A Randomized Controlled and Double-blinded Clinical Trial 99

Prapanbandit N

Effects of Kinesiotaping Combined with Medial Arch Support on Reduction of Heel Pain from Plantar Fasciitis: A Randomized Controlled Trial and Single-blinded 107

Ujaruspun N and Assawapalangchai S

Editorial Page

Functional Evaluation for Planning Exercise Programs in Patients with Chronic Diseases

Chronic diseases such as chronic obstructive pulmonary disease (COPD), chronic kidney disease, stroke, chronic pain, usually lead to impaired muscle strength and endurance, and affect movement functions. Therefore it is a role of physiatrists to perform functional evaluation so that appropriate rehabilitation programs could be planned.

In this issue, there are seven interesting original articles. Two studies used a 6-minute walk test (6MWT) and a 6-minute walk distance (6MWD) to evaluate cardiac and pulmonary functions: one used it to detect cardiac arrhythmia, an indicator to terminate phase II cardiac rehabilitation program and the other used it to detect risk of COPD exacerbation. The 6MWT is a practical evaluation that physiatrists could perform to evaluate patients' functions or fitness.

To restore muscle and movement functions, one needs exercise. High resistance and low repetition for strengthening muscle whereas low resistance and high repetition for improving muscle endurance. Weights and exercise equipment are commonly used for exercise trainings. However, these equipment may not be available and suitable for elderly. In this issue, one study introduced using a simple Thai-style braided rubber rope for a home-based exercise. It showed some improvement in muscle strength and quality of life of patients on continuous ambulatory peritoneal dialysis. However, patient's adherence to exercise is one of key success factors.

Another study in this issue is about poststroke depression which is prevalent during the first two years after stroke. Depression, if not well treated, affects motivation. Being a physiatrist, one should be aware of depression as it is a barrier to achieve optimal rehabilitation outcomes. Therefore one should be able to early detect depression which is rather common among those with chronic diseases seen in medical rehabilitation practice.

Apichana Kovindha, MD, FRCPhysiatrT
Editor